A Computer Program to Assess Egg Production Per Recruit in a Lobster *(Homarus americanus)* Population

G.P. Ennis and P.W. Collins

Fisheries Research Branch
Department of Fisheries and Oceans
P.O. Box 5667
St. John's, Newfoundland A1C 5X1

September 1983

Canadian Technical Report of Fisheries and Aquatic Sciences No. 1198

Canadian Technical Report of Fisheries and Aquatic Sciences

These reports contain scientific and technical information that represents an important contribution to existing knowledge but which for some reason may not be appropriate for primary scientific (i.e. *Journal*) publication. Technical Reports are directed primarily towards a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries management, technology and development, ocean sciences, and aquatic environments relevant to Canada.

Technical Reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report will be abstracted in *Aquatic Sciences and Fisheries Abstracts* and will be indexed annually in the Department's index to scientific and technical publications.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Details on the availability of Technical Reports in hard copy may be obtained from the issuing establishment indicated on the front cover.

Rapport technique canadien des sciences halieutiques et aquatiques

Ces rapports contiennent des renseignements scientifiques et techniques qui constituent une contribution importante aux connaissances actuelles mais qui, pour une raison ou pour une autre, ne semblent pas appropriés pour la publication dans un journal scientifique. Il n'y a aucune restriction quant au sujet, de fait, la série reflète la vaste gamme des intérêts et des politiques du Ministère des Pêches et des Océans, notamment gestion des pêches, techniques et développement, sciences océaniques et environnements aquatiques, au Canada.

Les Rapports techniques peuvent être considérés comme des publications complètes. Le titre exact paraîtra au haut du résumé de chaque rapport, qui sera publié dans la revue *Aquatic Sciences and Fisheries Abstracts* et qui figurera dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1-456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457-714, à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715-924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, Ministère des Pêches et de l'Environnement. Le nom de la série a été modifié à partir du numéro 925.

La page couverture porte le nom de l'établissement auteur où l'on peut se procurer les rapports sous couverture cartonnée.

Canadian Technical Report of
Fisheries and Aquatic Sciences 1198

September 1983

A COMPUTER PROGRAM TO ASSESS EGG PRODUCTION PER RECRUIT IN A LOBSTER (*HOMARUS AMERICANUS*) POPULATION

by

G. P. Ennis and P. W. Collins

Fisheries Research Branch
Department of Fisheries and Oceans
P.O. Box 5667
St. John's, Newfoundland A1C 5X1

This is the seventy-sixth Technical Report from
Fisheries Research Branch, St. John's, Newfoundland.

(c)Minister of Supply and Services Canada 1983

Cat. No. Fs 97-6/1198 ISSN 0706-6457

Correct citation for this publication:

Ennis, G. P., and P. W. Collins. 1983. A computer program to assess egg production per recruit in a lobster (Homarus americanus) population. Can. Tech. Rep. Fish. Aquat. Sci. 1198: iv + 15 p.

CONTENTS

Abstract/Résumé..... iv

Introduction..... 1

Biological basis of the analysis..... 1

Description of the program..... 4

Application of the program..... 4

References..... 4

Appendix 1: Input file layout and explanation of parameters..... 7

Appendix 2: Program listing..... 9

Appendix 3: Sample of program output..... 15

ABSTRACT

Ennis, G. P., and P. W. Collins. 1983. A computer program to assess egg production per recruit in a lobster (Homarus americanus) population. Can. Tech. Rep. Fish. Aquat. Sci. 1198: iv + 15 p.

A computer program which estimates the numbers of eggs produced per female recruit in a lobster population is described. The layout of the input file and an explanation of the parameters, a listing of the program and a sample of the output are provided as appendices. The program is run with various combinations of recruitment length (minimum legal size) and fishing mortality (exploitation rate) thereby providing a basis for assessing the impact of changes in size limit and/or exploitation rate in a lobster fishery on the production of eggs in the population.

Key words: lobster, egg production per recruit, assessment

RÉSUMÉ

Ennis, G. P., and P. W. Collins. 1983. A computer program to assess egg production per recruit in a lobster (Homarus americanus) population. Can. Tech. Rep. Fish. Aquat. Sci. 1198: iv + 15 p.

Le présent rapport décrit un programme informatisé pour l'estimation du nombre d'oeufs produits par femelle dans une population de homards. Le topogramme du fichier d'entrée, une explication des paramètres, un listage du programme et un état imprimé type constituent les annexes. Le programme est exécuté avec diverses combinaisons de longueurs au recrutement (taille légale minimum) et de mortalités par pêche (taux d'exploitation), générant ainsi une base pour l'évaluation de l'incidence des changement de la limite de taille ou du taux d'exploitation au cours de la pêche du homard sur la production d'oeufs dans la population.

INTRODUCTION

The basic management tools commonly employed in lobster (genus Homarus) fisheries are a minimum legal landing size (size limit) and effort limitation as a means of controlling exploitation rate. In theory at least, MSY can be realized from a population of lobsters if such regulatory measures are properly applied. However, it appears that the situation in most lobster fisheries can be represented by a point well to the right of MSY on a general production curve (i.e current landings are well below MSY and fishing effort is well in excess of that at MSY).

There are two facets to achieving MSY from a population. One is to maximize yield per recruit, the other is to maximize recruitment. Lobster fisheries are characterized by high exploitation rates and assessments have clearly demonstrated that, with existing size limits, yield per recruit is substantially less than maximum (Anon. 1977, 1979; Anthony and Caddy 1980). In addition, existing size limits are generally below the size at which most females lay eggs for the first time and with the excessively high exploitation rates that prevail in most areas, widespread recruitment overfishing is a distinct possibililty. In fact, recruitment overfishing appears to be the cause of stock collapse in certain areas of Eastern Canada (Robinson 1979).

Stock-recruitment relationships as such are poorly known for the genus Homarus, however, since current levels of landings are well below historical levels in most fisheries, it is reasonable to assume that, within the limits of habitat carrying capacity, increased egg production will result in increased recruitment. While it is clear that increasing the minimum legal size and/or reducing exploitation rates will result in greater numbers of eggs produced, it is only in very recent years that the detailed biological information required to properly assess the impact of changes in these fishery regulatory measures on egg production within a population have become available.

The purpose of this paper is to describe a computer program which has been written in order to carry out egg production per recruit analyses using the kind of information being obtained from studies of lobster population biology in Newfoundland waters.

BIOLOGICAL BASIS OF THE ANALYSIS

The biological concepts on which this egg per recruit program is based are very similar to those developed by Caddy (1979) in his relative population fecundity analysis. In the two analyses, however, there are different approaches to the way mortalities and growth are handled. Also, in the program presented here, there is provision to account for egg loss over and above that due to normal attrition as well as for those females which molt and extrude eggs in the same molting/spawning season.

The analysis is carried out as a chronological sequence of annual life history and mortality events which begins subsequent to the annual spring

fishing season and just prior to the annual molting/spawning period in summer. Initially, 1000 non-ovigerous females are distributed evenly at 1 mm size intervals over the 71-80 mm carapace length range and the sequence of life history and mortality events is repeated until these have disappeared from the population as a result of fishing or natural mortality. The analysis begins by estimating the number of individuals at each 1 mm CL that will extrude eggs during the molting/spawning period in year one. This estimate is obtained using a proportion derived from a functional size-maturity relationship (logisitc equation relating % mature and CL) which is based on detailed examination of pleopod cement glands done prior to the molting/spawning period as a means of determining whether individual animals will extrude eggs that year (Ennis, in press[1]). This relationship is represented by a sigmoid curve which sometimes does not fit the data very well at the upper and lower ends of the size range. In this analysis 100% maturity is assigned to those sizes at and above which the data indicate this to be the case, instead of assigning a lower percentage derived from the equation.

Some female lobsters molt prior to extruding eggs during the same annual molting/spawning period (Aiken and Waddy 1976, 1980). In annual (since 1975) shell condition sampling done at Arnold's Cove, Newfoundland following the molting period, the number of ovigerous females with new shells has ranged from 6.5 to 38.5% of the total number of ovigerous specimens examined (Ennis 1980). These have generally been smaller animals within the overall size range of ovigerous specimens and it was suggested that those females which molt and extrude eggs in the same year are spawning for the first time. These data have been examined in some detail and it has been found that the incidence of molting and spawning in the same year declines with increasing size. The data were subjected to probit analysis and a good fit obtained ($P > .99$). The resulting equation ($Y = 27.751 - 0.3072X$) is used to determine the number of those females which will extrude eggs in year one that molt before doing so. The number of eggs produced by these females is determined from the size-fecundity relationship using their postmolt length.

Estimates of proportions molting amongst those females which do not extrude eggs in year one are obtained from a relationship derived from molt predictions based on pleopod examination early in the molting/spawning period for non-ovigerous females but excluding those females whose pleopod cement glands indicate egg extrusion to be imminent. A quadratic regression was fitted to such data obtained from sampling at Arnold's Cove during June 9 to July 15, 1983 and the following equation obtained:

$$Y = -330.3593 + 13.5685X - 0.1073X^2 \text{ with } R^2 = .99.$$

The postmolt sizes of lobsters that molt are determined from a premolt-postmolt carapace length relationship obtained from sphyrion tagging (Ennis 1972, 1978).

No reliable estimates of natural mortality in lobsters are available but the general consensus reached by the ICES Working Group on Homarus Stocks is that it can be expected to be less than 10% annually for lobsters at sizes around the existing minimum legal sizes (Anon. 1977). One would expect most

natural mortality in lobsters to be associated with molting and in this analysis a 10% natural mortality rate is applied only to those lobsters that molt. Lobsters not molting in a particular year are subjected to a 5% rate of natural mortality.

Sphyrion tagging at Arnold's Cove in summer 1981 revealed that around 15% of the females which extrude eggs loose practically all of them over the incubation period (Ennis, in press[2]). Accordingly, the numbers of females which were estimated would extrude eggs were reduced by 15% to account for this egg loss. This percentage was applied uniformly with respect to size. Normal attrition of eggs over the 9-12 month incubation period has been estimated at around 36% (Perkins 1971). This egg loss is accounted for in the analysis by virtue of the fact that the size-fecundity relationship used to estimate the numbers of eggs produced is based on egg counts for ovigerous females taken near the end of the incubation period (Ennis 1981).

In addition to attrition of eggs over the incubation period, some female lobsters molt shortly after extruding thereby loosing the entire clutch of eggs (Ennis, in press[2]). This egg loss is accounted for in the functional size-maturity relationship. In a tag-recapture validation of egg extrusion predictions (based on cement gland staging), the only incorrect predictions recorded were for 2 out of 9 (22%) females with stage 2 cement glands (for which egg extrusion was predicted) which were not egg-bearing (but had molted) when recaptured prior to the following molting/spawning period. It is assumed that these two lobsters extruded eggs as predicted and lost them when they molted soon afterwards. Accordingly, in the data on which the functional size-maturity relationship is based, 22% of the animals with stage 2 cement glands were considered immature (Ennis, in press[1]).

So far the analysis has proceeded to the spring of year two and an estimate derived of the number of eggs extruded in year one that remain in the population and from which larvae will hatch during the summer. Except for those at the 71 and 72 mm carapace length intervals, most lobsters that molted in year one would have grown to or beyond the minimum legal size (81 mm CL) and at this stage in the analysis, except for those that laid eggs shortly after molting or were lost as a result of natural mortality, are subjected to fishing mortality. Egg-bearing females are protected from exploitation. In this analysis those females (15%) which loose their eggs over the incubation period are protected from exploitation as well because the presence of even very small numbers of eggs would make them illegal. All of these egg-bearing females (including those that loose their eggs) molt during the molting/spawning period in year two, are subjected to 10% natural mortality, and the survivors are subjected to fishing mortality during the fishing season in year three. For all other females remaining following the fishing season in year two, the initial stage in the analysis (i.e determining numbers that will extrude eggs during the upcoming molting/ spawning season) is now repeated and followed by the subsequent steps in the analysis.

The foregoing sequences are repeated until the 1000 individuals with which the analysis started have been removed from the population. The numbers of eggs produced and carried through the incubation period are then totalled.

DESCRIPTION OF THE PROGRAM

The program is written in Fortran IV and is very similar to the one developed by Ennis and Akenhead (1978) to assess yield per recruit in Newfoundland lobsters. Each component of the annual cycle of life history and mortality events is handled separately and in chronological order. Provision is made for different natural mortality rates for molters and non-molters, for the protection of egg-bearers from fishing mortality, for different growth rates for egg-bearers and non-egg-bearers, and for egg loss over the incubation period in excess of that due to normal attrition. In addition, special treatment is provided for females that molt and lay eggs in the same year.

The organization of the program is shown schematically in Fig. 1. The layout of the input file and an explanation of the parameters are provided in Appendix 1 and Appendix 2 is a listing of the program.

APPLICATION OF THE PROGRAM

The program is run with various combinations of recruitment length (minimum legal size) and fishing mortality (exploitation rate) thereby providing a basis for assessing the impact of changes in size limit and/or exploitation rate in a lobster fishery on the production of eggs in the population. The output (sample provided as Appendix 3) includes an estimate of the number of eggs carried to hatching produced by the 1000 females with which each analysis begins over the period that they remain in the population with a given combination of size limit and exploitation rate. Also included is an estimate of the numbers of eggs produced by these females at sizes smaller than the minimum legal size being considered.

REFERENCES

Aiken, D. E., and S. L. Waddy. 1976. Controlling growth and reproduction in the American lobster. p. 415-430. Proc. 7th Annual Meet. World Mariculture Soc.

1980. Reproductive Biology. Chapter 4. In The biology and management of lobsters, Vol. 1, Physiology and Behavior. J. S. Cobb and B. F. Phillips [ed.] Academic Press, 463 p.

Anon. 1977. Report of the Working Group on Homarus stocks. ICES C.M. 1977/K:11, 19 p.

1979. Report of the Homarus Working Group. ICES C.M. 1979/K:8, 49 p.

Anthony, V. C., and J. F. Caddy [eds.] 1980. Proceedings of the Canada-U.S. Workshop on Status of Assessment Science for N.W. Atlantic Lobster (Homarus americanus) Stocks (St. Andrews, N.B., Oct. 24-26, 1978). Can. Tech. Rep. Fish. Aquat. Sci. 932: vii + 186 p.

Caddy, J. F. 1979. Notes on a more generalized yield per recruit analysis for crustaceans, using size-specific inputs. Can. Fish. Mar. Serv. MS Rep. 1525: 7 p.

Ennis, G. P. 1972. Growth per moult of tagged lobsters (Homarus americanus) in Bonavista Bay, Newfoundland. J. Fish. Res. Board Can. 29: 143-148.

1978. Growth curves for Newfoundland lobsters from data on molt increment and proportion molting. CAFSAC Res. Doc. 78/29, 11 p.

1980. Size-maturity relationships and related observations in Newfoundland populations of the lobster (Homarus americanus). Can. J. Fish. Aquat. Sci. 37: 945-956.

1981. Fecundity of the American lobster, Homarus americanus, in Newfoundland waters. Fish. Bull. 79: 796-800.

Comparison of physiological and functional size-maturity relationships in two Newfoundland populations of lobsters, Homarus americanus. Fish. Bull. (in press[1]).

Tag-recapture validation of molt and egg extrusion predictions based upon pleopod examination in the American lobster, Homarus americanus. Fish. Bull. (in press[2]).

Ennis, G. P., and S. A. Akenhead. 1978. A model and computer program used to assess yield per recruit in Newfoundland lobster stocks. CAFSAC Res. Doc. 78/30, 13 p.

Robinson, D. G. 1979. Consideration of the lobster (Homarus americanus) recruitment overfishing hypothesis; with special reference to the Canso Causeway. p. 77-99. In F. D. McCracken [ed.] Canso Marine Environment Workshop. Part 3 of 4 Parts, Fishery Impacts. Can. Fish. Mar. Serv. Tech. Rep. 834: 163 p.

Perkins, H. C. 1971. Egg loss during incubation from offshore northern lobsters (Decapoda: Homaridae). Fish. Bull. 69: 451-453.



Fig. 1. Diagram showing organization of the program.

APPENDIX 1: INPUT FILE LAYOUT AND EXPLANATION OF PARAMETERS

RECORD LENGTH: 80 columns - card image

Card No.	Parameters	Format
1	LIST (20)	20A4
2	SLEN, FLEN, AMORT, BMORT CMORT, EGLOS, AFEC, BFEC AGR, BGR, IRUN	2F3.0, 4F5.0 4F10.0, I3
3	GDIST (10)	10F8.0
4	FSTRT, FSTOP, FINCR, RECLEN (5)	8F8.0
5-12	PROLAY (80)	10F8.0
13-20	PROMLT (80)	10F8.0
21-28	GROLAY (80)	10F8.0
29-36	PINIT (80)	10F8.0

1) LIST - Header of input file.

2) SLEN - Starting carapace length (mm).
FLEN - Last carapace length (mm).
AMORT - Proportion of molting component surviving natural mortality.
BMORT - Proporiton of laying component surviving natural mortality.
CMORT - Proporiton of old shells (i.e. those which neither molt not lay) surviving natural mortality.
EGLOS - Proportion of successful layers that eventually lose all eggs before hatching.
AFEC - Intercept of length-fecundity regression ($\log_{10}$).
BFEC - Slope of length-fecundity regression ($\log_{10}$).
AGR - Intercept of premolt-postmolt regression (arithmetic).
BGR - Slope of premolt-postmolt regression (arithmetic).
IRUN - Number of years (cycles).

3) GDIST - Distribution of growth scatter around postmolt mean length, derived from deviation around growth regression.

4) FSTRT - Starting exploitation rate.
FSTOP - Final exploitation rate.
FINCR - Increment of change for exploitation rate.
RECLEN - Recruit carapace lengths (mm) (maximum of five for each run).

5-12) PROLAY - Proportion of population laying eggs each length, derived from logistic equation relating % mature and CL based on pleopod cement gland development (see Ennis, In press).

13-20) PROMLT - Proportion of nonlaying component molting each length, derived from quadratic equation relating % molting amongst non-ovigerous females not going to lay egs and CL.

21-28) GROLAY - Proportion of laying component that also molt in same year each length, derived from probit equation based on incidencce of new-shelled ovigerous females in relation to size from fall sampling.

29-36) PINIT - Initial population numbers each length (start with 100 each length 71-80 mm).

```
FORTRAN IV G LEVEL  21                    MAIN

C     ***************************************************************
C     LOBSTER EGG PER RECRUIT PROGRAM
C     ***************************************************************
0001        INTEGER LIST(20),IGROW(80) NUMLEN,IRUN,ICNTR
0002        REAL RECLEN(5),POP(80)/80*0.0/,PROLAY(80),PROMLT(80)
0003        REAL CATCH(80),FEC(80),PINIT(80)/80*0.0/,GDIST(10)/10*0./
0004        REAL AMORT,BMORT,CMORT,EGLOS,AFEC,BFEC,AGR,BGR
0005        REAL FSTRT,FSTOP,FINCR,STARTN,NONLAY(80),NEGGS(80),PREG(80)
0006        REAL HATCHS(80),LOSERS(80),LOST(80),NOMOLT(80),NBOTH(80)
0007        REAL HOLD(80),MOLTER(80),POPLAY(80)/80*0.0/,GROLAY(80)
0008        REAL JUSLAY(80)
C                     ************************
C
C     SKIP REINITIALIZATION--FIRST TIME THRU
0009        GO TO 304
C
C     REINITIALIZE STORAGE FOR NEXT DATASET
0010    300     DO 301 I=1,NUMLEN
0011            POP(I)=0.
0012            PINIT(I)=0.
0013            HATCHS(I)=0.
0014            POPLAY(I)=0.
0015            NBOTH(I)=0.
0016            JUSLAY(I)=0.
0017            NEGGS(I)=0.
0018            PREG(I)=0.
0019            NONLAY(I)=0.
0020            LOSERS(I)=0.
0021            LOST(I)=0.
0022    301     NOMOLT(I)=0.
0023        DO 302 I=1,10
0024    302     GDIST(I)=0.
0025        DO 303 I=1,2
0026    303     RECLEN(I)=0.
0027    304 CONTINUE
C
C     READ TITLE---'D' IN COLUMN 80 GIVES DETAILED OUTPUT
0028        READ(5,2,END=305)LIST
0029        GO TO 306
0030    305 STOP
0031    306 CONTINUE
0032        WRITE(6,2)LIST
C
C     READ START LENGTH,LAST LENGTH,MOLTING SURVIVORSHIP,
C          LAYERS SURVIVORSHIP,OLD SHELL SURVIVORSHIP,
C          EGG-LOSS MORTALITY,INTERCEPT AND SLOPE OF FECUNDITY
C          REG.(LOG10),INTERCEPT AND SLOPE OF GROWTH REG.(ARITH),
C          NUMBER OF YEARS.
```

```
      C
0033        READ(5,1)SLEN,FLEN,AMORT,BMORT,CMORT,EGLOS,AFEC,BFEC,
           +AGR,BGR,IRUN
0034        NUMLEN=FLEN-SLEN
0035        NUMLEN=NUMLEN+1
0036        LS=SLEN
0037        LF=FLEN
0038        WRITE(6,13)LS,NUMLEN,AMORT,BMORT,CMORT,EGLOS,
           +AFEC,BFEC,AGR,BGR,IRUN
0039        LS=LS-1
      C
      C     READ PROPORTION OF SCATTER AROUND POST-MOLT LENGTH
0040        READ(5,5)GDIST
0041        WRITE(6,14)GDIST
      C
      C     READ VARIABLES OF EXPLOITATION RATE AND RECRUIT LENGTH
0042        READ(5,6)FSTRT,FSTOP,FINCR,RECLEN
0043        WRITE(6,15)FSTRT,FSTOP,FINCR,RECLEN
      C
      C     READ PROPORTION LAYERS EACH LENGTH
0044        READ(5,5) (PROLAY(I),I=1,NUMLEN)
      C
      C     READ PROPORTION MOLTING EACH LENGTH
0045        READ(5,5) (PROMLT(I),I=1,NUMLEN)
      C
      C     READ PROPORTION MOLTING AND LAYING SAME YEAR EACH LENGTH
0046        READ(5,5) (GROLAY(I),I=1,NUMLEN)
      C
      C     READ INITIAL POPULATION DISTRIBUTION
0047        READ(5,5) (PINIT(I),I=1,NUMLEN)
0048     20 CONTINUE
      C
      C     CALCULATE VECTOR OF FECUNDITY FOR LENGTHS
      C     PRINT HEADER FOR INPUT DATA
0049        WRITE(6,4)
0050          AFEC=10.**AFEC
0051          STARTN=0.
0052        DO 11 I=1,NUMLEN
0053          FEC(I)=AFEC*FLOAT(LS+I)**BFEC
0054            IGROW(I)=AGR+BGR*FLOAT(LS+I)
0055            IL=LS+I
0056            WRITE(6,3)IL,PINIT(I),PROMLT(I),PROLAY(I),GROLAY(I),
           +IGROW(I),FEC(I)
0057            STARTN=STARTN+PINIT(I)
0058     11 CONTINUE
      C
      C     ALL DATA NOW INPUT
      C     SET UP FISHING RATE LOOP AND CHECK FOR MORE VALUES
```

```
C       IF NONE RETURN FOR NEW DATASET AND REINITIALIZE
0059          ICNTR=0
0060    198   CONTINUE
0061          ICNTR=ICNTR+1
0062            IF(ICNTR.GT.5) GOTO 300
0063            IF(RECLEN(ICNTR).EQ.0.)GOTO 300
0064            LR=IFIX(RECLEN(ICNTR))
0065            FMORT=FSTRT-FINCR
0066    200     FMORT=FMORT+FINCR
0067            IF(FMORT.GT.FSTOP)GOTO 198
C
C       SET UP FOR LOOP THRU YEARS
0068            TEGGS=1.E-10
0069            TNUM=0.
0070            TCAT=0.
0071            TPREG=1.E-10
0072            THAT=0.
0073              DO 22 I=1,NUMLEN
0074                 HATCHS(I)=0.
0075                 POPLAY(I)=0.
0076                 JUSLAY(I)=0.
0077                 NBOTH(I)=0.
0078                 NEGGS(I)=0.
0079                 PREG(I)=0.
0080                 NONLAY(I)=0.
0081                 LOSERS(I)=0.
0082                 LOST(I)=0.
0083                 MOLTER(I)=0.
0084                 NOMOLT(I)=0.
0085    22           POP(I)=RINIT(I)
0086          WRITE(6,16)FMORT,LR
0087          IF(LIST(20).NE.    0)GOTO 23
0088          WRITE(6,18)
0089    23    CONTINUE
C
C       LOOP THRU IRUN NUMBER OF YEARS
0090          DO 100 IY=1,IRUN
C
C           SET UP FOR LOOP THRU LENGTHS
0091            SEGGS=0.
0092            SUMN=0.
0093            CSUM=0.
0094            SPREG=0.
0095            SHOLD=0.
0096            SHAT=0.
C
C           LOOP THRU LENGTHS
0097            DO 30 IK=1,NUMLEN
```

```
0098                    I=(NUMLEN-IK)+1
                C
                C       CALCULATE LAYERS,MOLTERS AND NON-MOLTERS
0099                    POPLAY(I)=POP(I)*PROLAY(I)
0100                    NONLAY(I)=POP(I)-POPLAY(I)
0101                    MOLTER(I)=NONLAY(I)*PROMLT(I)
0102                    NOMOLT(I)=NONLAY(I)-MOLTER(I)
                C
                C       CALCULATE NATURAL MORTALITY OF LAYERS,NON-MOLTERS
0103                    POPLAY(I)=POPLAY(I)*BMORT
0104                    NOMOLT(I)=NOMOLT(I)*CMORT
                C
                C       SEPARATE LAYING COMPONENT INTO LAYERS AND MOLTER/LAYERS
0105                    IF(GROLAY(I).EQ.0)GO TO 27
0106                    NBOTH(I)=POPLAY(I)*GROLAY(I)
0107                    JUSLAY(I)=POPLAY(I)-NBOTH(I)
0108                    NBOTH(I)=NBOTH(I)*AMORT
0109                    POPLAY(I)=0
                C
                C       DISTRIBUTE MOLTER/LAYERS AROUND PREDICTED POST MOLT
                C       MEAN LENGTH
0110                    JZ=(IGROW(I)-LS)-6
0111                       DO 26 K=1,10
0112                       JX=JZ+K
                C
                C          CHECK IF LEGAL SIZE TO GROW INTO
0113                       IF(JX.LT.1)JX=1
0114                       IF(JX.GT.NUMLEN)JX=NUMLEN
0115                       POPLAY(JX)=POPLAY(JX)+GDIST(K)*NBOTH(I)
0116                    26 CONTINUE
0117                    POPLAY(I)=POPLAY(I)+JUSLAY(I)
0118                 27 CONTINUE
                C
                C       CALCULATE NATURAL MORTALITY OF MOLTERS
                C       (I.E.  SURV. EGG-LOSERS+LAST YEARS LAYERS
                C       (THIS YEARS HATCHERS)+THIS YEARS MOLTERS
0119                    MOLTER(I)=MOLTER(I)+LOST(I)+HATCHS(I)
0120                    MOLTER(I)=MOLTER(I)*AMORT
                C
0121                    POP(I)=0.
                C
                C       DISTRIBUTE GROWERS AROUND PREDICTED POSTMOLT MEAN LENGTH
0122                    IZ=(IGROW(I)-LS)-6
0123                       DO 32 K=1,10
0124                          IX=IZ+K
                C
                C             CHECK THAT THIS IS A LEGAL SIZE TO GROW INTO
0125                          IF(IX.LT.1)IX=1
```

```
0126                        IF(IX.GT.NUMLEN)IX=NUMLEN
0127                        POP(IX)=POP(IX)+GDIST(K)*MOLTER(I)
0128                     CONTINUE
0129                  32 POP(I)=POP(I)+NOMOLT(I)
0130               30 CONTINUE
                   C
                   C     CALCULATE TOTAL FECUNDITY AND CATCH
0131                     DO 50 IJ=1,NUMLEN
0132                        I=(NUMLEN-IJ)+1
0133                        LOSERS(I)=HATCHS(I)*EGLOS
0134                        HATCHS(I)=HATCHS(I)-LOSERS(I)
0135                        NEGGS(I)=HATCHS(I)*FEC(I)
0136                        HOLD(I)=HATCHS(I)
0137                        PREG(I)=NEGGS(I)
0138                        IF(LS+I.GE.LR)PREG(I)=0.
0139                        CATCH(I)=POP(I)*FMORT
0140                        IF(LS+I.LT.LR)CATCH(I)=0.
0141                        HATCHS(I)=POPLAY(I)
0142                        LOST(I)=LOSERS(I)
0143                        POP(I)=POP(I)-CATCH(I)
0144                  50    CONTINUE
                   C
                   C     CALCULATE YEARLY SUMMATIONS
0145                     DO 31 I=1,NUMLEN
0146                        SEGGS=SEGGS+NEGGS(I)
0147                        SHOLD=SHOLD+HOLD(I)
0148                        SUMN=SUMN+POP(I)
0149                        CSUM=CSUM+CATCH(I)
0150                        SPREG=SPREG+PREG(I)
0151                        SHAT=SHAT+HATCHS(I)
0152               31 CONTINUE
                   C     IF DETAILED OPTION PRINT YEARLY SUMMATIONS
0153                     IF(LIST(20).NE.    0)GOTO 51
0154                     WRITE(6,19)SUMN,CSUM,SHAT,SHOLD,SEGGS
0155               51 CONTINUE
                   C
                   C     CALCULATE FINAL SUMMATIONS
0156                     TEGGS=TEGGS+SEGGS
0157                     TNUM=TNUM+SUMN
0158                     TCAT=TCAT+CSUM
0159                     TPREG=TPREG+SPREG
0160                     THAT=THAT+SHOLD
                   C
                   C     CHECK IF ANOTHER YEAR WILL CHANGE VALUES
0161                     IF(SUMN.LT.STARTN*.001)GOTO 101
0162              100 CONTINUE
0163                  WRITE(6,17)
0164               101 CONTINUE
```

```
C
C       CALCULATE YIELD VALUES
0165            YN=TEGGS/STARTN
0166            PP=TPREG/TEGGS
0167            PR=1.-PP
0168            WRITE(6,8)INUM,TCAT,THAT,TEGGS,TPREG,YN,PR,PP
C
C       RETURN FOR NEW VALUES OF EXPLOITATION RATE AND RECRUIT LENGTH
C
0169            GO TO 200
C       ****************************************************************
C       ********************FORMAT STATEMENTS*******************************
0170          1 FORMAT(2F3.0,4F5.0,4F10.0,I3)
0171          2 FORMAT(20A4)
0172          3 FORMAT(4X,I4,5X,F10.0,F12.4,5X,F12.4,5X,F12.4,8X,I10,8X,F9.0)
0173          4 FORMAT('0LENGTH(MM)     INIT POP.     PROP MOLTING     PROP LAYING
             *    PROP MOLT/LAY    POSTMOLT LENGTH     FECUNDITY')
0174          5 FORMAT(10F8.0)
0175          6 FORMAT(8F8.0)
0176          8 FORMAT(//' POPULATION NUMBERS:',F10.2/
             +' TOTAL CATCH:',5X,F10.2/
             +' TOTAL OVIGS:',5X,F10.2/
             +' TOTAL EGGS:',5X,F10.0/
             +' PRERECRUIT EGGS:',3X,F10.0/
             +' EGG YIELD/RECRUIT:',F10.0/
             +' PROP. RECRUIT EGGS:',F10.4/
             +' PROP. PRERECRUIT EGGS:',F10.4)
0177         13 FORMAT('0START LENGTH:',I5/'0NO. OF LENGTHS:',
             +I5/'0MOLTING SURVIVORSHIP:',F8.3/'0EGG-LAYING SURVIVORSHIP:',
             +F8.3/'0NON-MOLTING/LAYING SURVIVORSHIP:',F8.3/
             +'0EGG-LOSS MORTALITY:',F8.3/'0A AND B OF LOG10(FEC)=',
             +'A+B*LOG10(LENGTH):',2F10.5/'0A AND B OF POSTMOLT=',
             +'A+B*PREMOLT LENGTH:',2F10.5/'0NO. OF YEARS:',I5)
0178         14 FORMAT('0PROP. OF GROWERS AROUND PREDICTED LENGTH:',
             +/10X,10F8.4)
0179         15 FORMAT('0VARIABLE VALUES FOR EXPLOITATION RATE AND RECRUIT',
             +'LENGTH:'/10X,'FISH. MORT.: START   STOP  INCREMENT:',3F10.2/
             +' RECRUIT LENGTHS:',5F8.0)
0180         16 FORMAT(//' EXPLOITATION RATE:',F6.2,5X,'RECRUIT LENGTH:',I5/)
0181         17 FORMAT(' ***PROGRAM WENT BEYOND ALLOTED NUMBER OF YEARS*** ')
0182         18 FORMAT('0POPULATION          CATCH          OVIGS       HATCHED',
             +'     NO. OF EGGS')
0183         19 FORMAT(' ',3X,F6.2,8X,F6.2,7X,F6.2,6X,F6.2,4X,F10.0)
C       ****************************************************************
0184            END
```

APPENDIX 3: SAMPLE OF PROGRAM OUTPUT

Exploitation rate: 0.80 Recruit length: 81

Population numbers:	509.79
Total catch:	893.07
Total ovigs:	434.71
Total eggs:	5076735.
Prerecruit eggs:	2463943.
Egg yield/recruit:	5077.
Prop. recruit eggs:	0.5147
Prop. prerecruit eggs:	0.4853